

口腔殺菌消毒劑
舒立克喉嚨噴液劑
THROATEC Spray for Throat

衛部藥製字第 059628 號
醫師藥師藥劑生指示藥品

【有效成分及含量】每毫升含POVIDONE IODINE 4.5mg

【其他成分(賦形劑)】GLYCERIN、PROPYLENE GLYCOL、ETHANOL、POTASSIUM IODIDE、L-MENTHOL、PURIFIED WATER。

【用途(適應症)】口腔消毒殺菌。

【使用上注意事項】

- 有下列情形者，請勿使用：
 - (1)曾因本藥成分引起過敏的人。
 - (2)對碘過敏的人。
- 有下列情形者，使用前請洽醫生診治：
 - (1)孕婦、可能懷孕婦女及哺乳婦。
- 有下列情形者，使用前請先諮詢醫師藥師藥劑生：
 - (1)口內有潰爛的人。
 - (2)有甲狀腺疾病的人。
- 其他使用上注意事項：
 - (1)存放於小孩不會隨手拿到的地方。
 - (2)密栓保管於日光不會直射並且涼爽的地方。
 - (3)請依建議用法用量。
 - (4)不要長期連續使用。
 - (5)本藥限於口腔使用，不得內服。
 - (6)注意不要噴入眼睛，萬一跑入眼睛要馬上用清水洗，馬上接受眼科的診察。
 - (7)會使皮膚或衣物染色，沾到衣服上時要儘快用水或洗劑清洗。
 - (8)避免誤用。為保持品質，不要換裝到其他容器。
 - (9)不要靠近火源的地方。
 - (10)超過保存期限後，請勿使用。
 - (11)不要用針等物品，扎弄噴嘴前端的洞。
 - (12)在冬季或氣溫低的時候，中間的粘度會較高，會有噴射較不容易的情形，請放在手中稍微加溫後再使用。
 - (13)小孩使用時要在大人的協助下使用。
 - (14)假牙或其他含銀材質之牙科材料可能會因本劑而變色。
 - (15)有全身性吸收之可能性，因此反覆使用新生兒可能出現甲狀腺功能低下，孕婦及其胎兒及哺乳婦有造成甲狀腺腫及甲狀腺功能低下。
 - (16)會使皮膚或衣物染色。

【用法用量】

用 法	用 量
1.打開瓶蓋。 2.將噴嘴朝向喉嚨尖部直接使用。 3.使用後蓋緊瓶蓋。 先噴4~5次，直到第一次有液體	一天數次，適量使用於喉嚨粘膜處。

【警語】

1.使用本藥後，若有發生以下副作用，請立即停止使用，並持此說明書諮詢醫師藥師藥劑生。

關係部位	症 狀
消化器官	不快感，噁心，嘔吐，口內潰爛感染，灼熱感，刺激感
皮 膚	紅潮，蕁麻疹，發疹
呼吸系統	呼吸困難
其 他	胸口悶，浮腫

2.使用本藥後，若有發生以下症狀時，請立即停止使用，並接受醫師診治：

- (1)若有任何不適情況產生。
- (2)自覺狀況沒有改善或更惡化。
- (3)服用本藥4~5天後，仍不見症狀改善。
- (4)使用後出現蕁麻疹、浮腫、胸悶等症狀及臉色蒼白、手脚冰冷、冒冷汗、呼吸困難等現象。

【包裝】4000毫升以下塑膠瓶裝

【儲存】25℃以下儲存



委託者
BEST PHARMA CO. LTD.
倍斯特醫藥生物科技股份有限公司
總公司：台中市北屯區崇德路二段51號3樓之3、之5
製造廠：利達製藥股份有限公司
廠 址：台中市大甲區中山路一段906號

Oral Antiseptic Spray
THROATEC® Spray for Throat

【Active ingredient and content】 POVIDONE IODINE 4.5mg/ml

【Other ingredients (Excipient)】 GLYCERIN · PROPYLENE GLYCOL · ETHANOL · POTASSIUM IODIDE · L-MENTHOL · PURIFIED WATER.

【Use (Indication)】 Spray into throat or onto affected area with several sprays per day for sore throat.

【Precautions】

1. Do not use for people having the following conditions:

- (1) Who experienced an allergic reaction.
- (2) Who is allergic to iodine.

2. Please consult a doctor before use:

- (1) Who are pregnant, wish to get pregnant, or are lactating (breastfeeding).

3. Please consult a doctor or pharmacist:

- (1) Who are aphthous ulcer.
- (2) Who have thyroid diseases.

4. Other precautions:

- (1) Keep out of reach of children.
- (2) Seal and store in a cool place away from direct sunlight.
- (3) Rigid adherence to the recommendation.
- (4) Not recommended for long term use.
- (5) For external use only.
- (6) In case get into the eyes, flush eyes with water immediately and consult your doctor as soon as possible.
- (7) Wash thoroughly with water or detergent once stained clothing.
- (8) Do not transfer to other containers to avoid improper use and to maintain quality.
- (9) Keep from contact with fire.
- (10) Use before the expiry date.
- (11) Do not prick the hole in the front part of the spray nozzle.
- (12) Keep warm in hands before use in winter time or low temperature.
- (13) For children use should be supervised by adults.
- (14) Could be bled on your false toothedenture.
- (15) Possibility of systemic absorption, and during breastfeeding increases breastmilk iodine levels and can cause transient hypothyroidism in breastfed infants.
- (16) Could dye on skin or clothing.

【Administration and dosage】

Administration	Dosage
1. Open bottle cap. 2. Spray into throat or affected area directly. 3. Cap tightly after use. Spray 4~5 times till the liquid is spouted.	Spray into throat or onto affected area with several sprays per day.

【Warnings】

1. If you notice any side effects, please consult a doctor or pharmacist with this insert.

Affected areas	Symptoms
Digestive organs	unpleasant feeling, nausea, vomiting, apotheosis, burning, irritating
Skin	redness, urticarial, rashes
Expiratory	Dyspnea (breathing difficulties)
Others	Chest distress, edema

2. If you notice any side effects, please stop to use and consult a doctor immediately:

- (1) If any discomfort occurs.
- (2) If your symptoms don't improve or worsen.
- (3) After use 4~5 days, still no improvement in symptoms.
- (4) Appear after use, urticaria, edema, chest tightness, pale, cold hands and feet, cold sweat or difficulty breathing.

【Package】 Plastic bottle under 4000 ml

【Storage】 Store below 25°C



Under the authority of
BEST PHARMA CO. LTD.
3F.-3-5 No. 51, SEC 2 Chung Der Road,
Tai-Chung, Taiwan, 406, R.O.C

BMB-003